

Traffic light system for identifying risk of serious illness

	Green – low risk	Amber – intermediate risk	Red – high risk
Colour (of skin, lips or tongue)	<ul style="list-style-type: none"> • Normal colour 	<ul style="list-style-type: none"> • Pallor reported by parent/carer 	<ul style="list-style-type: none"> • Pale/mottled/ashen/blue
Activity	<ul style="list-style-type: none"> • Responds normally to social cues • Content/smiles • Stays awake or awakens quickly • Strong normal cry/not crying 	<ul style="list-style-type: none"> • Not responding normally to social cues • No smile • Wakes only with prolonged stimulation • Decreased activity 	<ul style="list-style-type: none"> • No response to social cues • Appears ill to a healthcare professional • Does not wake or if roused does not stay awake • Weak, high-pitched or continuous cry
Respiratory		<ul style="list-style-type: none"> • Nasal flaring • Tachypnoea: <ul style="list-style-type: none"> – RR >50 breaths/minute, age 6–12 months – RR >40 breaths/minute, age >12 months • Oxygen saturation ≤95% in air • Crackles in the chest 	<ul style="list-style-type: none"> • Grunting • Tachypnoea: RR >60 breaths/minute • Moderate or severe chest indrawing
Circulation and hydration	<ul style="list-style-type: none"> • Normal skin and eyes • Moist mucous membranes 	<ul style="list-style-type: none"> • Tachycardia: <ul style="list-style-type: none"> – >160 beats/minute, age <12 months – >150 beats/minute, age 12–24 months – >140 beats/minute, age 2–5 years • CRT ≥3 seconds • Dry mucous membranes • Poor feeding in infants • Reduced urine output 	<ul style="list-style-type: none"> • Reduced skin turgor
Other	<ul style="list-style-type: none"> • None of the amber or red symptoms or signs 	<ul style="list-style-type: none"> • Age 3–6 months, temperature ≥39°C • Fever for ≥5 days • Rigors • Swelling of a limb or joint • Non-weight bearing limb/not using an extremity 	<ul style="list-style-type: none"> • Age <3 months, temperature ≥38°C* • Non-blanching rash • Bulging fontanelle • Neck stiffness • Status epilepticus • Focal neurological signs • Focal seizures
CRT, capillary refill time; RR, respiratory rate * Some vaccinations have been found to induce fever in children aged under 3 months			
This traffic light table should be used in conjunction with the recommendations in the NICE guideline on fever in under 5s.			