

Best Practice Guidelines for the Management and Transfer of Patients with a diagnosis of Ruptured Abdominal Aortic Aneurysm to a specialist Vascular Centre



Endorsed by



The College of
Emergency Medicine



Summary of recommendations

1. A clinical diagnosis of ruptured abdominal aortic aneurysm (rAAA) should be considered:
 - In patients over the age of 50 years presenting with abdominal/back pain AND hypotension
 - In patients with a known AAA and symptoms of either abdominal/back pain OR hypotension/collapse
 - In patients where an alternative diagnosis is considered more likely on clinical grounds, rAAA still must be excluded, with radiological confirmation made prior to referral.
Level 3, strong recommendation
2. Permissive hypotension is advocated for patients with a clinical diagnosis of rAAA to maintain an alert patient and systolic blood pressure >70mm Hg is acceptable. *Level 4, strong*
3. If a specialist vascular service cannot be provided on-site the patient requires transfer to a centre with appropriate facilities and expertise. Transfer agreements with the local ambulance service should be in place. *Level 4, strong recommendation*
4. Rapid and co-ordinated transfer can reduce delays in the patient journey and improve outcome. *Level 3, strong.* To expedite transfer the most senior doctor available should lead and be actively involved the care of any patient with suspected rAAA. Outgoing referrals should go to a senior vascular trainee or consultant.

Items 5–18 below are all Level 5, with strong recommendation
5. All patients with a clinical or radiological diagnosis of rAAA should be assessed as to their current clinical state AND pre-morbid level of function to determine suitability for transfer.
6. Patients aged <85 with no/mild/moderate systemic disease should be referred to the receiving hospital's on-call vascular service without delay.
7. Patients age >85 or with severe systemic disease will benefit from a consultant*–consultant discussion prior to transfer to a vascular unit.
8. Impaired mental capacity is not a contraindication to assessment and transfer.
9. Patients who have been previously turned down for elective surgery should still be discussed via a consultant*–consultant referral.
10. Contraindications to transfer are restricted to those with cardiac arrest in the current admission and intubated patients. Such patients are unlikely to survive transfer and surgery.
11. There are no ESSENTIAL investigations required prior to transfer. However, a blood gas and an emergency department ultrasound are considered useful, if these incur no delay.

12. Investigations including FBC, U&E, amylase, X-match, CT scans MUST not delay transfer to a centre that can provide definitive care. If an alternative diagnosis is more likely, or the investigation can be performed without causing delay, it is reasonable to perform these investigations before transfer.
13. Patients should be treated, if necessary, with both analgesia (according to the College of Emergency Medicine [CEM] guidelines) and fluids before and during transfer. Blood products and inotropes may be required, their use should be supervised by a ST4 or above, trainee or equivalent or consultant.
14. A time-critical transfer in a 999 ambulance, preferably with a paramedic crew is required, although this is not essential.
15. Patients requiring inotropic support will need a suitably experienced and trained medical escort for transfer.
16. The facility to transfer CT images electronically must be in place to ensure all images are transferred to the receiving hospital. If electronic transfer is not possible, a CD or DVD of DICOM files must accompany the patient.
17. Patients should not travel with blood products, unless transfusion already commenced.
18. Patients who remain cardiovascularly stable should be transferred to either an emergency department resuscitation bed or local equivalent. Patients who are unstable may need rapid transfer to theatres.
19. Transfer to a specialist vascular centre should occur within 30 minutes of diagnosis.

**if consultants are off-site, out of hours, the most senior person involved in patient care should initiate the discussion/referral*

Scope

This guideline aims to ensure patients with a clinical diagnosis of rAAA equity of access to a specialist vascular centre for expert assessment and intervention. If not already present, emergency physicians and vascular surgeons should work collaboratively to develop region-wide protocols to facilitate access to this standard of care. Simultaneously this guideline should assist the emergency physician in the assessment, resuscitation and timely preparation for transfer of the patient with a clinical diagnosis of ruptured abdominal aortic aneurysm, if this diagnosis is made in a hospital without suitable vascular facilities.

This guideline should optimise and standardise the management of patients arriving in vascular centres; ensuring that patients who would benefit from surgery are transferred without delay and ensure those not suitable for surgery, and likely to die, are palliated appropriately.

Reason for development

There are no existing UK guidelines on the emergency management of rAAA, although these are available for other critical conditions and life-threatening conditions; eg, head injury, asthma. There is evidence of wide variation in current practice in the emergency management of patients with rAAA (see below).

There is an accelerating trend for vascular and endovascular surgery to be practised in fewer, but larger, vascular centres, which is likely to increase the need for the transfer of patients from presenting centres to centres where definitive treatment can be offered.¹⁻³ The views of a cohort of vascular specialists (including radiologists) and emergency physicians have been obtained, using the Delphi consensus method.⁴

Variation in practice

Currently, there appears to be widespread variation in practice, including the selection of patients for referral, diagnosis, referral pathways and the conditions under which the transfer is made⁴ which may incur unnecessary time delays between presentation and repair.

Introduction/background

rAAA is a life-threatening emergency. Without rapid surgical or endovascular intervention, there is an almost 100% mortality within 3 days.⁵ With appropriate intervention mortality can be as low as 20%. Surgical intervention is most effective if delivered rapidly, reducing the duration of hypotension that leads to multi-organ failure in the post-operative period on ICU. Although hypotension may be fluid responsive, aggressive pre-operative fluid therapy may accelerate bleeding and permissive pre-operative hypotension is recommended. There is some evidence, from retrospective single centre cohort studies, that an untimely or delayed transfer may worsen patient outcome.⁶⁻⁸ Any unnecessary delay can prolong the period of hypotension, worsening eventual outcome. Therefore, delivery of the patient to a specialist who can provide definitive care is time critical. National 30-day mortality rates for

repair of rAAA are >40%,⁹ but mortality results of <20% have been reported by a few specialist centres outside of the UK.^{10,11}

Management

Emergency management of rAAA in a hospital without a specialist vascular centre has six steps.

1. Diagnosis
2. Assessment of suitability for transfer
3. Referral to a vascular specialist
4. Monitoring and treatment in the emergency department
5. Transfer to the vascular specialist
6. Further management by the vascular specialist (repair and subsequent inpatient care).

1 Diagnosis

Symptoms/ signs of rAAA include abdominal/back pain, collapse with pallor, hypotension (which initially may be postural only), tachycardia, sweating, agitation (usually secondary to pain or hypotension), pulsatile abdominal mass, pain radiating to the legs and evidence of lower limb ischaemia.¹² In elderly patients, flank pain should prompt consideration of rAAA and patients may be misdiagnosed with renal colic. Flank pain/peri-umbilical discoloration (Cullens/ Grey-Turners) are late signs of rAAA.

Clinical diagnosis

A clinical diagnosis of ruptured abdominal aortic aneurysm can be made:

- In patients over the age of 50 years presenting with acute onset abdominal/ back pain AND hypotension
- In patients with a known AAA symptoms of either abdominal/ back pain OR hypotension/ collapse.

Patients with other combinations of signs/ symptoms may need further investigation to confirm/ exclude the diagnosis of rAAA.

Radiological diagnosis

Ultrasound can confirm the presence and size of an aortic aneurysm.¹³ It cannot exclude rupture. However, ultrasound confirmation of an aortic aneurysm in a patient with abdominal/ back pain or hypotension supports the clinical diagnosis of rAAA, and a well-visualised, normal aorta in a haemodynamically stable patient excludes rAAA. Ultrasound has the advantage of portability, it does not require contrast and the complications thereof and its increasing availability in the emergency

department as a result of an increasing number of trained emergency physicians skilled in its use. The disadvantages of ultrasound include the fact it is operator-dependent, adequate visualization of the aorta may be impaired by body habitus or bowel gas and a rupture cannot be confirmed. A further disadvantage is that a normal-sized aorta viewed on ultrasound in the presence of severe hypotension does not exclude the diagnosis of rAAA or other abdominal pelvic aneurysm, in which case a contrast-enhanced diagnostic CT scan must be performed without delay.

CT scan can confirm the diagnosis of rAAA and is recommended in stable patients pre-operatively to confirm diagnosis and assist operative planning.^{10,14} If image acquisition is to specific protocol, CT provides rapid high resolution images which can be reconstructed quickly in any plane allowing for rapid endovascular planning or the identification of adverse features for both endovascular and open repair. It might also provide alternative/co-morbid diagnoses. As there are some associated disadvantages; the requirement for image transfer, interpretation by a radiologist, contrast requirement in patients who might have impaired renal function and the potential need for repeat scans if the diagnostic scan is of insufficient quality to plan endovascular repair or unreadable in the receiving hospital. The CT scan may be best performed at the vascular centre if the clinical diagnosis is secure without CT scan.

It is essential that electronic data transfer between referring hospitals in a region is in place. For imaging this requires all regional hospitals in England to transfer images via the image exchange portal (IEP) but in the other 3 UK countries more sophisticated bespoke systems already exist and must be used. Facilities for cross border image transfer must be in place. In the absence of reliable electronic transfer systems, image transfer should be by transfer of DICOM files to CD or DVD.

2. Assess suitability for transfer

Physiological and pre-morbid condition is more important than age in assessing suitability for transfer of a patient to a specialist vascular centre. Blood pressure and conscious level are the most important physiological parameters.^{10,11}

The extent of pre-existing systemic disease should be assessed as none, mild, moderate or severe.

Identify whether the patient has lost consciousness in the current episode.¹⁵

Identify whether there is evidence (from notes or relatives) that the patient has given an advance directive.

Patients with cardiac arrest in the current episode have poor outcomes and are not candidates for transfer.¹⁶

3 Referral to a vascular specialist

Speed of referral and transfer is critical once the likely diagnosis has been made and confirmed by an experienced emergency medicine doctor. For alert patients of 85 years or less, with no, mild or moderate systemic disease this request can be made by an experienced emergency medicine doctor to a

senior vascular trainee in the closest specialist vascular centre. Under most circumstances, the patient does not need to be assessed by a local general surgeon prior to the request for transfer, although if no experienced emergency medicine doctor is available, the patient may need to be assessed by a local general surgeon. For candidates over 85 years, or with severe systemic disease, who are intubated or have fluctuating consciousness, the possibility of referral (or decision not to transfer) must be made after discussion between the most senior emergency doctor available and the on-call vascular surgeon at the closest vascular centre.

4 Monitoring and treatment in the emergency department

The patient's vital signs must be monitored continuously.

Investigations

If the patient is known to have an abdominal aortic aneurysm, the clinical diagnosis does not have to be supported by investigations.

Investigations should not delay the transfer of patients with a strong clinical suspicion of aneurysm rupture but the following investigations should be considered:

- Abdominal ultrasound or CT scan
- ECG
- Full blood count, serum electrolytes and creatinine, serum amylase, troponin, blood gas.

Treatments

There are no essential treatments required prior to or during transfer, that are known to affect outcome. A timely decision regarding transfer, as discussed above, and then prompt transfer if required are the key therapeutic interventions. Management in the emergency department should focus on alleviating pain and maintaining adequate perfusion, whilst preparing the patient for transfer (and theatre). On occasion critical care interventions may be required.

Pain can be managed according to standard CEM guidance.¹⁷

Large bore intravenous access should be obtained. Permissive hypotension is advocated and fluid resuscitation to maintain the patient mentally alert with a systolic BP >70mm Hg is acceptable (titrated to a systolic of 70–90 mmHg with cautious crystalloids).^{10,18} Blood products may be required, but cross matching must not delay transfer. Rarely inotropic support may be required if the patient is not fluid responsive.

Patients who are or have been unconscious will require intubation and assisted ventilation prior to transfer.

Patients should be kept nil by mouth. Insertion of a urinary catheter and/or central venous line should only occur if they do not delay transfer. All further preparations for theatre can be undertaken at the receiving centre.

5 Transfer to a specialist vascular centre

Patients should be transferred swiftly by an emergency ambulance. A paramedic crew is preferred but not essential. A trained transfer doctor escort only is required for patients requiring inotropic support or with a blood transfusion already commenced.

Any cross-sectional imaging performed locally may aid the receiving vascular unit, to expedite aneurysm repair. However, the organisation of image transfer, either electronically or as downloaded DICOM files, should not be allowed to incur delay. Similarly patients should not under normal circumstances travel with cross-matched blood or other blood products,²¹ unless they have been commenced prior to transfer.

After agreement to accept the patient, the vascular senior registrar/consultant should put in place all necessary arrangements to assess and manage the patient expeditiously upon arrival.

Patients should be transferred to an area which offers critical care facilities (monitored bed, oxygen, direct nursing care, doctor availability). The exact location will depend on local arrangements but is likely to include Emergency Department resuscitation bed or critical care (ICU / high dependency). Some (particularly unstable) patients may be best transferred directly to the operating theatre or intervention suite.

6 Further care by vascular specialist

This is beyond the scope of this guideline but there is some evidence that patient outcomes improve with increasing hospital volume^{1,3} and the availability and use of formal protocols for patient care.^{19,20}

Audit standard

Transfer to specialist care should occur within 30 minutes of diagnosis.

References

- 1 Holt PJ, Karthikesalingam A, Poloniecki JD, et al. Propensity scored analysis of outcomes after ruptured abdominal aortic aneurysm. *Br J Surg* 2010;97:496-503.
- 2 Wyatt M, Lamont P, Naylor R, et al. The Provision of Services for Patients with Vascular Disease 2012. www.vascularsociety.org.uk
- 3 Holt PJ, Poloniecki JD, Gerrard D et al. Meta-analysis and systematic review of the relationship between volume and outcome in abdominal aortic aneurysm surgery. *Br J Surg* 2007; 94: 395-403.
- 4 Hinchliffe RJ, Ribbons TLP, Ulug P, Powell JT. The transfer of patients with ruptured abdominal aortic aneurysm from general hospitals to specialist vascular centres: results of a Delphi consensus study. Submitted to *Emerg Med J* February 2012.
- 5 Choksy SA, Wilmink AB, Quick CR. Ruptured abdominal aortic aneurysm in the Huntingdon district: a 10-year experience. *Ann R Coll Surg Engl* 1999;81:27-31.
- 6 Hames H, Forbes TL, Harris JR et al. The effect of patient transfer on outcomes after rupture of an abdominal aortic aneurysm. *Can J Surg* 2007;50:43-7.
- 7 Haveman JW, Karlicsek A, Verhoeven EL et al. Results of a streamlined regional ambulance transport and subsequent treatment of acute abdominal aortic aneurysm. *Emerg Med J* 2006;23:807-10.
- 8 Salhab, Farmer J, Osman I. Impact of delay on survival in patients with ruptured abdominal aortic aneurysm. *Vascular* 2006;14:38-42.
- 9 Bown MJ, Sutton AJ, Bell PR, Sayers RD. A meta-analysis of 50 years of ruptured abdominal aortic aneurysm repair. *Br J Surg*. 2002 Jun;89:714-30.
- 10 Mayer D, Pfammatter, Rancic Z et al. 10 years of emergency endovascular aneurysm repair for ruptured abdominal aortoiliac aneurysms: lessons learned. *Ann Surg* 2009;249:510-5.
- 11 Opfermann P, von Allmen R, Diehm N et al. Repair of ruptured abdominal aortic aneurysm in octogenarians. *Eur J Vasc Endovasc Surg*. 2011;42:475-83.
- 12 Basnyat PS, Biffin AH, Moseley LG, et al. Mortality from ruptured abdominal aortic aneurysm in Wales. *Br J Surg*. 1999;**86**:765-70.

- 13 Dent B, Kendall RJ, Boyle AA, Atkinson PR. Emergency ultrasound of the abdominal aorta by UK emergency physicians: a prospective cohort study. *Emerg Med J* 2007;24:547-9.
- 14 Moll FL, Powell JT, Fraedrich G et al. Management of abdominal aortic aneurysms clinical practice guidelines of the European Society for Vascular Surgery. *Eur J Vasc Endovasc Surg* 2011;41:S1-58.
- 15 Acosta S, Ogren M, Bergqvist D et al. The Hardman index in patients operated on for ruptured abdominal aortic aneurysm: a systematic review. *J Vasc Surg* 2006;44:949-54.
- 16 Greeven AP, Bouwman LH, Smeets HJ et al. Outcome of patients with ruptured abdominal aortic aneurysm after cardiopulmonary resuscitation. *Acta Chir Belg* 2011;111:78-82.
- 17 Management of pain in adults. 2010. www.collemergencymed.ac.uk
- 18 van der Vliet JA, van Aalst DL, Schultze LJ et al. Hypotensive hemostasis (permissive hypotension) for ruptured abdominal aortic aneurysm: are we really in control? *Vascular* 2007;15:197-200.
- 19 Moore R, Nutley M, Cina CS et al. Improved survival after implementation of an emergency endovascular therapy protocol for ruptured abdominal aortic aneurysms. *J Vasc Surg* 2007;45:443-50.
- 20 Davies MG, Younes HK, Harris PW et al. Outcomes before and after initiation of an acute aortic treatment center. *J Vasc Surg* 2010;52:1478-85.
- 21 UK Blood Transfusion Services, Transfusion Handbook, 2007 4th Editions, www.transfusionguidelines.org.uk