

USE OF GLYCERYL TRINITRATE (GTN)

Ref No: 3678

Lead Clinician	:	Dr Adrian Marsh, Consultant Emergency Medicine
Business Centre	:	Emergency & Critical Care Centre
Developed (date)	:	December 2012
Last updated	:	May 2013
Last reviewed	:	May 2013
Planned review	:	December 2014
Hyperlink (where applicable)	:	
Keywords	:	GTN, Glyceryl Trinitrate
Comments	:	

INDICATIONS

Acute pulmonary oedema

Uncontrolled pain of cardiac origin, including aortic dissection

Accelerated hypertension with pulmonary oedema or acute coronary syndrome

DOSAGE

Acute pulmonary oedema: glyceryl trinitrate (GTN) by continuous IV infusion, initially 20 microgram/min, increasing in increments of 20 microgram/min at 15–30 min intervals until desired response or a maximum of 200 microgram/min is achieved, provided BP remains >90/60 mmHg

Uncontrolled pain of cardiac origin: GTN by continuous IV infusion, initially 10 microgram/min, titrated upwards at 15-min intervals in increments of 5 or 10 microgram/min according to patient response to a maximum of 200 microgram/min if necessary to control pain, provided BP remains >90/60 mmHg

Accelerated hypertension: follow dosage instructions according to clinical presentation (see above); otherwise give GTN by continuous IV infusion, initially 5 microgram/min, titrated upwards at 15-min intervals in increments of 5 or 10 microgram/min according to patient response at to a maximum of 100 microgram/min

PREPARATIONS

GTN 1 mg/mL in 50 mL vial

ADMINISTRATION

- Fill a compatible 50 mL syringe (see [Notes](#)) with GTN solution 1 mg/mL (50 mL)
- Administer via a syringe pump, and titrate according to patient response ([Table 1](#))

NOTES

Compatible syringes and tubing; rigid plastic syringes (e.g. Gillette Sabre, Brunswick Disposable, BD Plastipak); polyethylene tubing (e.g. Vygon Lectrocath, David Bull Laboratories Types A261 or A2001) or nitro extension set.

GTN is **incompatible** with polyvinylchloride (PVC) infusion bags (e.g. Steriflex, Boots, Viaflex, Travenol) and PVC tubing as up to 50% of the GTN can be lost onto the plastic requiring bigger doses.

Table 1: GTN infusion 1mg/ml via syringe pump (flow rate – mL/hr)

Dosage (microgram/min)	5	10	15	20	25	30	35	40	45	50
Flow rate (mL/hr)	0.3	0.6	0.9	1.2	1.5	1.8	2.1	2.4	2.7	3
Dosage (microgram/min)	55	60	65	70	75	80	85	90	95	100
Flow rate (mL/hr)	3.3	3.6	3.9	4.2	4.5	4.8	5.1	5.4	5.7	6
Dosage (microgram/min)	105	110	115	120	125	130	135	140	145	150
Flow rate (mL/hr)	6.3	6.6	6.9	7.2	7.5	7.8	8.1	8.4	8.7	9
Dosage (microgram/min)	155	160	165	170	175	180	185	190	195	200
Flow rate (mL/hr)	9.3	9.6	9.9	10.2	10.5	10.8	11.1	11.4	11.7	12

Patient Monitoring

The patient must be attached to a cardiac monitor and have continuous heart rate, minimum of 3 lead monitoring, oxygen saturations. Blood pressure should be measured every 5 minutes for the first 30 minutes, every 15 minutes for the next 30 minutes and then every 30 minutes. If the blood pressure drops to less than 90/60, the infusion should be stopped and an urgent medical review requested. The infusion can be restarted with the rate decreased by 10 microgram/min once the blood pressure is greater than 90/60 mmHg.