

# CLINICAL REVIEW

## Gallstones

Kurinchi S Gurusamy *lecturer*, Brian R Davidson *professor of hepatopancreatobiliary and liver transplantation surgery*

Department of Surgery, 9th floor Royal Free Hospital, Royal Free Campus, UCL Medical School, London, UK

Gallstones affect approximately 5-25% of adults in the Western world. It is therefore important to understand the consequences of a diagnosis of gallstones, the associated complications, and treatment to allow patients to be appropriately advised. The purpose of this review is to update clinicians on the diagnosis and management of gallstones.

### What are gallstones?

Gallstones are crystalline deposits in the gallbladder (figure 1).<sup>1</sup> The prevalence of gallstones varies between 5% and 25%, with a higher prevalence in Western countries, women, and older age group.<sup>2</sup> Traditionally, gallstones were classified as cholesterol stones, pigment stones, or mixed stones (a combination of cholesterol and pigment stones) based on their composition,<sup>3</sup> which can only be determined reliably after their removal.<sup>4</sup> Recently, additional types of gallstones have been identified based on their microscopic structure and composition.<sup>1</sup> However, most stones fall under the umbrella of cholesterol (37-86%), pigment (2-27%), calcium (1-17%), or mixed (4-16%).<sup>1-4</sup> The types of gallstone vary by their cause, the measures attempted to prevent their formation, their appearance on radiographs, and their response to dissolution therapy. The current recommendations for diagnosis and management are, however, the same for all types of gallstone.

### Who gets gallstones?

Cholesterol stones are formed because of the alteration in the balance between pronucleating factors and antinucleating factors in the bile. Factors that lead to gallstone formation include excessive bile cholesterol, low bile salt levels, decreased gallbladder motility, and the phosphatidylcholine molecule, which prevents the crystallisation of cholesterol.<sup>5</sup> The main risk factors for cholesterol stone formation include female sex, pregnancy, high dose oestrogen treatment, increasing age, ethnicity (higher prevalence in Native American Indians and lower prevalence in black Americans, Africans, and people from China, Japan, India, and Thailand), genetic traits, obesity, high serum triglyceride levels, low levels of high density cholesterol, rapid weight loss, high calorific diet, refined carbohydrate diet, lack of physical activity, cirrhosis, Crohn's disease, and gallbladder stasis (for example, as a result of previous

gastrectomy or vagotomy).<sup>6-8</sup> Haemolysis and chronic bacterial or parasitic infections are considered the main risk factors for pigment stones<sup>5</sup> and are preventable causes of gallstones.

### Can gallstone formation be prevented?

Although some of the causes of gallstones, such as obesity, rapid weight loss, a high calorific diet, a refined carbohydrate diet, and lack of physical activity are preventable by lifestyle changes, there is currently no evidence that lifestyle modifications can reduce the incidence of gallstones. Haemolysis and infections can be prevented by early recognition of sickle cell disease, taking appropriate measures for prevention of sickling crises, and using prophylactic antibiotics in those who have undergone splenectomy or had splenic infarction. Another way of preventing gallstone formation is to remove the gallbladder in people undergoing anti-obesity operations (as rapid weight loss is one of the risk factors for gallstone formation) and other major abdominal operations to avoid further surgery as a result of the development of symptomatic gallstones. There is currently no evidence to suggest that prophylactic cholecystectomy is indicated in any patient group without gallstones<sup>9</sup> or that any of the above suggested measures of preventing gallstones are effective.

### How do gallstones present?

Each year approximately 2-4% of people with gallstones develop symptoms, with biliary colic being the most common symptom (steady right upper quadrant abdominal pain lasting more than half an hour)<sup>10-14</sup> in the absence of fever. Presence of fever usually indicates acute cholecystitis or cholangitis. Other common symptoms related to gallstones include epigastric pain and intolerance to fried or fatty foods (symptoms such as nausea, bloating, flatulence, frothy and foul smelling stools).<sup>14</sup> The box lists the complications resulting from gallstones, and includes acute cholecystitis (0.3-0.4% annually),<sup>11-13 15</sup> acute pancreatitis (0.04-1.5% annually),<sup>15</sup> obstructive jaundice (0.1-0.4% annually),<sup>12 13 15</sup> and other rarer complications such as acute cholangitis and intestinal obstruction (gallstone ileus). Of these, acute pancreatitis and cholangitis are life threatening complications, with 3% to 20% mortality after a first attack of acute pancreatitis<sup>16</sup> and 24% mortality after acute cholangitis.<sup>17</sup>

### Summary points

- Gallstones are common in adults
- Currently, treatment is indicated only for symptomatic gallstones
- Pain is the most common symptom related to gallstones
- Occasionally gallstones present with life threatening complications
- Surgery is the only definitive treatment and is recommended for people with symptomatic gallstones who are fit to undergo surgery
- Laparoscopic cholecystectomy is currently the preferred method of surgery and is generally both effective and safe; however, there is a 0.3% risk of serious injury to the bile duct, which may have serious long term consequences

### Sources and selection criteria

We searched Medline, Embase, the Cochrane Database of Systematic Reviews, and Clinical Evidence online using the search terms "gallstones" or "cholelithiasis", focusing mainly on systematic reviews, meta-analyses, and high quality randomised controlled trials published within the past five years, wherever appropriate and possible.

Uncomplicated biliary colic often precedes other gallstone related complications.<sup>11</sup> The rates of gallstone related complications are higher in people with a history of uncomplicated biliary colic.<sup>18</sup> Although studies have shown an association between gallstones and cancer of the biliary tract,<sup>19 20</sup> no causative link has been established and the observed association could be due to the presence of common factors causing gallstones and gallbladder cancer.<sup>21</sup>

## How should suspected gallstones be investigated?

Ultrasonography is currently the first line method for the diagnosis of gallstones and has a high diagnostic accuracy (90% sensitivity and 88% specificity) even when performed by non-radiologists.<sup>22</sup> Based on the agreement in a consensus conference, the diagnosis of acute cholecystitis is suspected by the presence of local or systemic signs of inflammation, such as Murphy's sign (tenderness in the right upper quadrant below the costal margin on deep inspiration; sensitivity 65% and specificity 87%<sup>23</sup>), fever, increased white cell count or C reactive protein, and confirmed by ultrasonography, computed tomography, or magnetic resonance imaging.<sup>24</sup> Radiological signs of acute cholecystitis include a thickened gallbladder wall (>4 mm), an enlarged gallbladder (long axis diameter >8 cm, short axis diameter >4 cm), or fluid collection around the gallbladder.<sup>24</sup> The diagnosis of pancreatitis is usually suspected by the presence of pain in the epigastric region radiating to the back and confirmed by diffuse abdominal tenderness, increased serum amylase, urine amylase, or serum lipase levels, and is supported by radiological features such as an enlarged pancreas with peripancreatic fluid collections.<sup>25</sup> The consensus conference conducted by the European Association for Endoscopic Surgery concluded that common bile duct stones should be suspected by the presence of clinical features suggestive of obstructive jaundice, such as yellowish discoloration of skin and dark urine supported by an increased serum bilirubin or alkaline phosphatase level and confirmed with magnetic resonance cholangiopancreatography or endoscopic ultrasonography.<sup>26 27</sup> Fever and rigors in the presence of jaundice should raise the suspicion of cholangitis.

If patients present with symptoms suggestive of gallstones, are systemically well, and do not have features suggestive of acute cholecystitis, acute pancreatitis, obstructive jaundice, or cholangitis, it is reasonable to investigate them by an elective ultrasonography, followed by elective referral to a general surgeon if gallstones are present. If gallstone complications are suspected, urgent referral to the surgeon is warranted because

early confirmation of diagnosis and treatment of complications are associated with better outcomes (see section on timing of surgery). Features that suggest the presence of complications include fever, rigors, hypotension, epigastric pain radiating to the back, dark urine, jaundice, Murphy's sign, diffuse abdominal tenderness, or a positive result for urine bile pigments on urinalysis. Depending on the clinical presentation, further blood tests, such as blood white cell count, levels of serum C reactive protein, serum amylase, serum bilirubin, and serum alkaline phosphatase; urine tests to check levels of urine amylase and urine lipase; and radiological investigations such as ultrasonography, computed tomography, magnetic resonance imaging, magnetic resonance cholangiopancreatography, and endoscopic ultrasonography may be performed to confirm or rule out the presence of gallstones and complications.

## How is gallstone disease treated?

### Asymptomatic gallstones

The distinction between symptomatic and asymptomatic gallstones can be difficult as symptoms can be mild and varied. While gallstone complications can be diagnosed using one or more of the criteria described, in patients presenting with vague upper abdominal pain or dyspeptic symptoms it can be difficult to discern whether the symptoms are related to gallstones. In one study, 90% of patients with classic biliary colic had high rates of symptom relief after cholecystectomy, suggesting that biliary colic is a fairly reliable indicator of symptomatic gallstones.<sup>28</sup> Around 70% of patients with upper abdominal pain with no further restriction by intensity or duration of pain had symptom relief after cholecystectomy. Only 55% of patients with dyspeptic symptoms had symptom relief, suggesting that in a major proportion of people vague upper abdominal pain or dyspepsia may not be related to gallstones.<sup>28</sup> There is currently no evidence that lifestyle modifications such as decreasing fatty food intake or increasing exercise decreases or prevents the incidence of symptoms in people with asymptomatic gallstones. No treatment is currently recommended for patients with asymptomatic gallstones (irrespective of whether these are cholesterol, pigment, or mixed stones) except for patients with porcelain gallbladders (which is usually identified by ultrasonography), owing to the association with gallbladder cancer.<sup>29</sup> The reason for advising against surgery for asymptomatic gallstones is because of the complications associated with surgical intervention, although this is a topic of ongoing debate.

In patients with asymptomatic gallstones undergoing major abdominal surgery, it seems reasonable to offer

### Complications of gallstones

- Acute cholecystitis
- Choledocholithiasis
- Acute cholangitis
- Acute pancreatitis
- Mucocele of gallbladder
- Empyema of gallbladder
- Gangrenous gallbladder
- Biliary peritonitis
- Porcelain gallbladder
- Gallbladder cancer

cholecystectomy, as adhesions related to a major operation may make further minimal access surgeries difficult or impossible. There is, however, no evidence from randomised controlled trials or systematic reviews to support this statement.

### Symptomatic gallstones

Cholecystectomy (removal of the gallbladder) is the preferred option in the treatment of gallstones, irrespective of whether the gallstones are cholesterol, pigment, or mixed stones. Evidence from randomised controlled trials, systematic reviews, and cohort studies show that extracorporeal shock wave lithotripsy or bile acid dissolution therapy with ursodeoxycholic acid has a low rate of cure, with only 27% of patients having dissolution of stones after treatment with ursodeoxycholic acid and only 55% of carefully selected patients being stone free after extracorporeal shock wave lithotripsy. The rate of recurrent gallstones is also high; more than 40% of patients have recurrence of gallstones within four years after complete dissolution of stones or extracorporeal shock wave lithotripsy. Over three months, only 26% of people remained colic free after treatment with ursodeoxycholic acid compared with 33% after placebo, and about 2% of people had gallstone complications after treatment with ursodeoxycholic acid, which is similar to the annual rate of complications in those not taking the drug.<sup>15 30-34</sup> In patients who are not suitable for cholecystectomy because of their general medical condition, percutaneous cholecystostomy (temporary external drainage of the gallbladder contents through a tube inserted under radiological guidance) may be considered in an emergency situation, although a systematic review revealed that the role of percutaneous cholecystostomy in the management of such patients was not clear.<sup>35</sup> When the patient's condition has improved, cholecystectomy may be reconsidered. Based on evidence from randomised controlled trials, watchful observation may be a suitable alternative to surgery in a small proportion of people who do not get recurrent symptoms.<sup>36 37</sup> It is, however, not possible to predict those patients who will get recurrence of symptoms.

### *Cholecystectomy: the risks and benefits*

Although cholecystectomy is a relatively safe procedure with few serious complications, bile duct injury resulting from surgery is a serious complication, with potential long term consequences.<sup>38</sup> The overall short term mortality after surgery varies between 0% and 0.3%.<sup>9</sup> Although traditionally less than 0.5% of people undergoing cholecystectomy are believed to have bile duct injury,<sup>9</sup> a study of more than 50 000 unselected patients from the Swedish Registry for Gallstone Surgery and ERCP, GallRiks, revealed that 1.5% of patients undergoing cholecystectomy between 2005 and 2010 developed a bile duct injury, although only a fifth of these injuries (0.3%) involved

partial or complete transection of the bile duct.<sup>38</sup> The patients with bile duct injury had a significantly higher one year mortality compared with those without such an injury.<sup>38</sup>

Cholecystectomy is generally performed by key hole operation (laparoscopic cholecystectomy)<sup>39</sup> because of the shorter length of hospital stay, decreased pain, earlier return to work, and better cosmesis. Laparoscopic cholecystectomy can be performed as a day procedure and generally involves four incisions measuring less than 1 cm each. Fat intolerance may develop in a small proportion of people after cholecystectomy, and a low fat diet is recommended in these patients. However, there is currently no strong evidence to support the usefulness of such a diet.

In patients with symptomatic gallbladder stones and common bile duct stones, the treatment options include open cholecystectomy with open exploration of the common bile duct, laparoscopic cholecystectomy with laparoscopic exploration of the common bile duct, and laparoscopic cholecystectomy with endoscopic sphincterotomy (performed preoperatively, intraoperatively, or postoperatively).<sup>40</sup> Evidence from a systematic review of randomised controlled trials shows that there is no evidence of difference in the morbidity or incidence of retained stones between endoscopic sphincterotomy and laparoscopic exploration of the common bile duct and inconsistency as to whether there is any difference in the length of hospital stay between the two approaches.<sup>40</sup>

### *When is the optimum time for surgery?*

The timing of surgery for various indications is controversial. In patients with biliary colic there is no medical reason to delay surgery, the delays being caused only by the availability of resources (although surgery can be delayed by surgeons recommending weight reduction for particular patients). Evidence from a randomised controlled trial, which compared early surgery within 24 hours of hospital admission versus delayed surgery with an average wait of about four months on the waiting list, showed that delaying surgery increased complications (0% in early group versus 22.5% in the delayed group) and hospital stay (an average of one additional day).<sup>41</sup> The timing of cholecystectomy in patients with acute cholecystitis is also controversial. Although the traditional belief was to allow the inflammation to settle and perform laparoscopic cholecystectomy after a period of at least six weeks, a systematic review on this topic has shown that early laparoscopic cholecystectomy performed within one week of onset of symptoms can avoid further complications from gallstones while waiting for surgery.<sup>42</sup> Early laparoscopic cholecystectomy can also decrease hospital stay by about four days without increasing surgical complications (approximately 5-6% in each group) or the proportion of people requiring conversion from laparoscopic to open surgery (approximately 20% in each group).<sup>42</sup> Although

most of the gallbladder related complications during the waiting time in the delayed group in the studies included in the systematic review were recurrence or non-resolution of acute cholecystitis, there is potential for further episodes of pain, pancreatitis, or obstructive jaundice while waiting. Evidence from a randomised controlled trial showed that morbidity after laparoscopic cholecystectomy between seven and 45 days was approximately two or three times that of surgery performed early; hence surgery within this timeframe is not recommended.<sup>43</sup> For timing of surgery in patients with mild acute pancreatitis (no organ failure or local complications), evidence from a systematic review that included only one small randomised controlled trial showed that performing surgery as early as possible (rather than waiting until symptoms settle and for blood test results to return to normal levels) decreased the hospital stay by one day,<sup>44</sup> although experts have expressed concerns that the severity of pancreatitis may not be evident until 48 hours and surgery in patients with severe pancreatitis (organ failure or local complications) within 48 hours can be harmful.<sup>45</sup> Delaying surgery for 48 hours overcomes this concern. The two circumstances where early cholecystectomy may not be appropriate are in patients with severe acute pancreatitis and those presenting during pregnancy.<sup>9,44</sup> Further trials are necessary to resolve these problems.

## What is the impact of gallstone disease on health services and society?

In 2004 in the United States, a total of 1.8 million outpatient visits were related to gallstones.<sup>46</sup> Each year, more than 0.5 million cholecystectomies are performed in the United States<sup>47</sup> and 70 000 in England.<sup>48</sup> The cost of cholecystectomy and loss of working time because of symptoms related to gallstones, and their treatment has an important impact on health services and society.

The illustration was produced by Rakhee Bashar from UCL Medical Illustration in collaboration with the authors.

Contributors: KG performed the literature search, wrote the article, and is the guarantor. BRD critically commented on the article.

Competing interests: We have read and understood the BMJ Group policy on declaration of interests and declare the following interests: BRD acts as an expert witness in court proceedings related to gallstone management but this has no influence on the submitted work.

Provenance and peer review: Commissioned; externally peer reviewed.

- Qiao T, Ma RH, Luo XB, Yang LQ, Luo ZL, Zheng PM. The systematic classification of gallbladder stones. *PLoS One* 2013;8:e74887.
- Kratzer W, Mason RA, Kachele V. Prevalence of gallstones in sonographic surveys worldwide. *J Clin Ultrasound* 1999;27:1-7.
- Trotman BW, Soloway RD. Pigment vs cholesterol cholelithiasis: clinical and epidemiological aspects. *Am J Dig Dis* 1975;20:735-40.
- Schafmayer C, Hartleb J, Tepel J, Albers S, Freitag S, Volzke H, et al. Predictors of gallstone composition in 1025 symptomatic gallstones from Northern Germany. *BMC Gastroenterol* 2006;6:36.
- Van Erpecum KJ. Pathogenesis of cholesterol and pigment gallstones: an update. *Clin Res Hepatol Gastroenterol* 2011;35:281-7.
- Stinton LM, Myers RP, Shaffer EA. Epidemiology of gallstones. *Gastroenterol Clin North Am* 2010;39:157-69, vii.
- Shaffer EA. Epidemiology and risk factors for gallstone disease: has the paradigm changed in the 21st century? *Curr Gastroenterol Rep* 2005;7:132-40.
- Banim PJ, Luben RN, Bulluck H, Sharp SJ, Wareham NJ, Khaw KT, et al. The aetiology of symptomatic gallstones quantification of the effects of obesity, alcohol and serum lipids on risk. Epidemiological and biomarker data from a UK prospective cohort study (EPIC-Norfolk). *Eur J Gastroenterol Hepatol* 2011;23:733-40.
- Gurusamy KS, Davidson BR. Surgical treatment of gallstones. *Gastroenterol Clin North Am* 2010;39:229-44, viii.
- Berger MY, van der Velden JJM, Lijmer JG, de Kort H, Prins A, Bohnen AM. Abdominal symptoms: do they predict gallstones? A systematic review. *Scand J Gastroenterol* 2000;35:70-6.
- Attili AF, De Santis A, Capri R, Repice AM, Maselli S. The natural history of gallstones: the GREPCO experience. The GREPCO Group. *Hepatology* 1995;21:655-60.

- Del Favero G, Caroli A, Meggiato T, Volpi A, Scalon P, Puglisi A, et al. Natural history of gallstones in non-insulin-dependent diabetes mellitus. A prospective 5-year follow-up. *Dig Dis Sci* 1994;39:1704-7.
- Halldestam I, Enell EL, Kullman E, Borch K. Development of symptoms and complications in individuals with asymptomatic gallstones. *Br J Surg* 2004;91:734-8.
- Portincasa P, Moschetta A, Petruzzelli M, Palasciano G, Di Ciaula A, Pezzolla A. Gallstone disease: symptoms and diagnosis of gallbladder stones. *Best Pract Res Clin Gastroenterol* 2006;20:1017-29.
- Venneman NG, van Erpecum KJ. Gallstone disease: primary and secondary prevention. *Best Pract Res Clin Gastroenterol* 2006;20:1063-73.
- Yadav D, Lowenfels AB. Trends in the epidemiology of the first attack of acute pancreatitis: a systematic review. *Pancreas* 2006;33:323-30.
- Salek J, Livote E, Sideridis K, Bank S. Analysis of risk factors predictive of early mortality and urgent ERCP in acute cholangitis. *J Clin Gastroenterol* 2009;43:171-5.
- Rutledge D, Jones D, Rege R. Consequences of delay in surgical treatment of biliary disease. *Am J Surg* 2000;180:466-9.
- Hsing AW, Gao YT, Han TQ, Rashid A, Deng J, Chen J, et al. Gallstones and the risk of biliary tract cancer: a population-based study in China. *Br J Cancer* 2007;97:1577-82.
- Ishiguro S, Inoue M, Kurahashi N, Iwasaki M, Sasazuki S, Tsugane S. Risk factors of biliary tract cancer in a large-scale population-based cohort study in Japan (JPHC study); with special focus on cholelithiasis, body mass index, and their effect modification. *Cancer Causes Control* 2008;19:33-41.
- Shebl FM, Andreotti G, Meyer TE, Gao YT, Rashid A, Yu K, et al. Metabolic syndrome and insulin resistance in relation to biliary tract cancer and stone risks: a population-based study in Shanghai, China. *Br J Cancer* 2011;105:1424-9.
- Ross M, Brown M, McLaughlin K, Atkinson P, Thompson J, Powelson S, et al. Emergency physician-performed ultrasound to diagnose cholelithiasis: a systematic review. *Acad Emerg Med* 2011;18:227-35.
- Trowbridge RL, Rutkowski NK, Shojania KG. Does this patient have acute cholecystitis? *JAMA* 2003;289:80-6.
- Mayumi T, Takada T, Kawarada Y, Nimura Y, Yoshida M, Sekimoto M, et al. Results of the Tokyo consensus meeting Tokyo guidelines. *J Hepatobiliary Pancreat Surg* 2007;14:114-21.
- Sunamura M, Lozonchi L, Takeda K, Kobari M, Matsuno S. Criteria for diagnosis of acute pancreatitis in Japan and clinical implications. *Pancreas* 1998;16:243-9.
- Williams EJ, Green J, Beckingham I, Parks R, Martin D, Lombard M, et al. Guidelines on the management of common bile duct stones (CBDSt). *Gut* 2008;57:1004-21.
- Scientific Committee of the European Association for Endoscopic Surgery. Diagnosis and treatment of common bile duct stones (CBDSt). Results of a consensus development conference. *Surg Endosc* 1998;12:856-64.
- Berger MY, Olde Hartman TC, Bohnen AM. Abdominal symptoms: do they disappear after cholecystectomy? *Surg Endosc* 2003;17:1723-8.
- Gurusamy KS, Samraj K. Cholecystectomy versus no cholecystectomy in patients with silent gallstones. *Cochrane Database Syst Rev* 2007;1:CD006230.
- Strasberg SM, Clavien PA. Overview of therapeutic modalities for the treatment of gallstone diseases. *Am J Surg* 1993;165:420-6.
- Venneman NG, Besselink MG, Keulemans YC, Vanberge-Henegouwen GP, Boermeester MA, Broeders IA, et al. Ursodeoxycholic acid exerts no beneficial effect in patients with symptomatic gallstones awaiting cholecystectomy. *Hepatology* 2006;43:1276-83.
- May GR, Sutherland LR, Shaffer EA. Efficacy of bile acid therapy for gallstone dissolution: a meta-analysis of randomized trials. *Aliment Pharmacol Ther* 1993;7:139-48.
- Carrilho-Ribeiro L, Pinto-Correia A, Velosa J, Carneiro De Moura M. A ten-year prospective study on gallbladder stone recurrence after successful extracorporeal shock-wave lithotripsy. *Scand J Gastroenterol* 2006;41:338-42.
- O'Donnell LD, Heaton KW. Recurrence and re-recurrence of gall stones after medical dissolution: a longterm follow up. *Gut* 1988;29:655-8.
- Gurusamy KS, Rossi M, Davidson BR. Percutaneous cholecystostomy for high-risk surgical patients with acute calculous cholecystitis. *Cochrane Database Syst Rev* 2013;8:CD007088.
- Schmidt M, Sondenaa K, Vethrus M, Berhane T, Eide GE. A randomized controlled study of uncomplicated gallstone disease with a 14-year follow-up showed that operation was the preferred treatment. *Dig Surg* 2011;28:270-6.
- Schmidt M, Sondenaa K, Vethrus M, Berhane T, Eide GE. Long-term follow-up of a randomized controlled trial of observation versus surgery for acute cholecystitis: non-operative management is an option in some patients. *Scand J Gastroenterol* 2011;46:1257-62.
- Tornqvist B, Stromberg C, Persson G, Nilsson M. Effect of intended intraoperative cholangiography and early detection of bile duct injury on survival after cholecystectomy: population based cohort study. *BMJ* 2012;345:e6457.
- Department of Health. NHS reference costs 2012 to 2013. [www.gov.uk/government/publications/nhs-reference-costs-2012-to-2013](http://www.gov.uk/government/publications/nhs-reference-costs-2012-to-2013).
- Dasari BV, Tan CJ, Gurusamy KS, Martin DJ, Kirk G, McKie L, et al. Surgical versus endoscopic treatment of bile duct stones. *Cochrane Database Syst Rev* 2013;12:CD003327.
- Salman B, Yuksel O, Irkoculu O, Akurek N, Tezcaner T, Dogan I, et al. Urgent laparoscopic cholecystectomy is the best management for biliary colic. A prospective randomized study of 75 cases. *Dig Surg* 2005;22:95-9.
- Gurusamy KS, Davidson C, Gluud C, Davidson BR. Early versus delayed laparoscopic cholecystectomy for people with acute cholecystitis. *Cochrane Database Syst Rev* 2013;6:CD005440.
- Gutt CN, Encke J, Koninger J, Harnoss JC, Weigand K, Kipfmuller K, et al. Acute cholecystitis: early versus delayed cholecystectomy, a multicenter randomized trial (ACDC study, NCT00447304). *Ann Surg* 2013;258:385-93.
- Gurusamy KS, Nagendran M, Davidson BR. Early versus delayed laparoscopic cholecystectomy for acute gallstone pancreatitis. *Cochrane Database Syst Rev* 2013;9:CD010326.
- Bouwense SA, Bakker OJ, van Santvoort HC, Boerma D, van Ramshorst B, Gooszen HG, et al. Safety of cholecystectomy in the first 48 hours after admission for gallstone pancreatitis not yet proven. *Ann Surg* 2011;253:1053-4; author reply 54-5.
- Everhart J. The burden of digestive diseases in the United States. 2008. [www3.niddk.nih.gov/Burden\\_of\\_Digestive\\_Diseases/](http://www3.niddk.nih.gov/Burden_of_Digestive_Diseases/).
- Centers for Disease Control and Prevention. Discharges with at least one procedure in nonfederal short-stay hospitals, by sex, age, and selected procedures: United States, selected years 1990 through 2009-2010. [www.cdc.gov/nchs/hus/content2012.htm#098](http://www.cdc.gov/nchs/hus/content2012.htm#098).



**Questions for future research**

Can lifestyle modifications such as increased physical activity, a low fat diet, and avoidance of high calorific and refined sugar diet prevent or reduce the development of gallstones or can they decrease the incidence and severity of symptoms related to gallstones?

What is the optimal timing of cholecystectomy in patients with acute severe pancreatitis and pregnant women with symptomatic gallstones?

What is the optimal surgical technique to avoid complications related to laparoscopic cholecystectomy?

What is the optimal approach to reducing the symptoms of cholecystectomy and facilitating increased day case surgery and earlier return to work?

What new approaches could be used in patients with symptomatic gallstones who are unfit to undergo surgery?

**Tips for non-specialists**

If biliary colic is suspected, ultrasonography remains the first line investigation to establish whether gallstones are present


If acute cholecystitis or acute gallstone pancreatitis is suspected, emergency investigations and immediate referral to a specialist are recommended, as early surgery may be indicated

48 Health and Social Care Information Centre. Hospital episode statistics, Admitted patient care, England—2012-13. [www.hscic.gov.uk/catalogue/PUB12566/hosp-epis-stat-admi-proc-2012-13-tab.xlsx](http://www.hscic.gov.uk/catalogue/PUB12566/hosp-epis-stat-admi-proc-2012-13-tab.xlsx).

Cite this as: *BMJ* 2014;348:g2669

© BMJ Publishing Group Ltd 2014

## Figure



Obstruction to the common bile duct by common duct stones may cause jaundice or cholangitis. Obstruction to the pancreatic duct or ampulla of Vater by common duct stones may cause pancreatitis