

Utility of bedside leucocyte esterase testing to rule out septic arthritis

Thomas Knapper ¹, Richard J Murphy ¹, Brett Rocos,¹ James Fagg,² Nick Murray,³ Michael Richard Whitehouse ^{1,4}

Handling editor David Metcalfe

¹Department of Trauma and Orthopaedics, North Bristol NHS Trust, Westbury on Trym, UK

²Department of Trauma and Orthopaedics, Royal United Hospitals Bath NHS Foundation Trust, Bath, UK

³Trauma and Orthopaedics, Great Western Hospital, Swindon, UK

⁴Musculoskeletal Research Unit, University of Bristol School of Clinical Science, Bristol, UK

Correspondence to

Mr Thomas Knapper, Department of Trauma and Orthopaedics, North Bristol NHS Trust, Westbury on Trym, Bristol, BS10 5NB, UK; tdknapper@me.com

Received 1 May 2020

Revised 14 January 2021

Accepted 11 February 2021

Published Online First

10 March 2021

ABSTRACT

Introduction Suspected septic arthritis is a common presentation to EDs. The underlying diagnosis is often non-infective pathology. Differentiating between aetiologies is difficult. A bedside test with high negative predictive value (NPV) may allow safe discharge of patients, reduce the time in the ED, hospital admission and associated costs. This study aims to evaluate the NPV of bedside leucocyte esterase (LE) in the assessment of these patients.

Methods A prospective multicentre observational study of ED adult patients referred to orthopaedics with suspected native joint septic arthritis between October 2015 and April 2016. At three hospital sites in the Bristol region, the results of the LE test exposed to aspirated synovial fluid were recorded along with Gram stain, culture, haematinics and length of stay. A positive LE test was considered 2+ or 3+ leucocytes based on the test strip colour. Data were analysed to establish sensitivity, specificity, NPV and positive predictive value (PPV) against the gold standard 48-hour culture. We determined the potential number of inpatient bed-days that might be avoided using this bedside test.

Results Eighty patients underwent joint aspiration. Five cases had positive 48-hour culture. All (5/5) infected cases showed $\geq 2+$ LE, sensitivity of 100% (95% CI 47.8% to 100%) while the Gram stain was positive in only one case (sensitivity 20%, 95% CI 0.51% to 71.6%). Twenty-three LE were read negative or 1+, all with negative 48-hour culture results, resulting in an NPV of 100% (95% CI 82.1% to 1.00%) for a negative LE test. Specificity of a positive LE test was 30.7% (95% CI 20.5% to 42.45%) with PPV of 8.77% (95% CI 7.64% to 10.1%). It was calculated that 57 orthopaedic bed-days could have potentially been saved by immediately discharging those with a negative LE test.

Conclusions LE point-of-care testing for suspected septic arthritis of native joints has a high NPV. Implementation of LE may facilitate more rapid discharge of patients with negative results. This test has the potential to reduce diagnostic uncertainty and costs to the healthcare system.

INTRODUCTION

Bacterial septic arthritis is an orthopaedic emergency. In the adult population referrals to the orthopaedic team for suspected native joint infection (NJI) are common; however incidence is low, 7.8 per 100 000 in the UK.¹ The detection rate for NJI varies between published studies with a range of 8%–27%.^{2–4} Risk factors for septic arthritis include: extremes of age, diabetes mellitus, intravenous

Key messages

What is already known on this subject

- ▶ Studies have shown leucocyte esterase (dipstick) to have high sensitivity and specificity during the investigation of suspected prosthetic joint infection, and to be a quick and reliable test in the evaluation of pleural and peritoneal aspirates.
- ▶ Few studies have established the usefulness of leucocyte esterase in the investigation and exclusion of a native joint infection and those that exist have looked at predominantly paediatric or young adult populations, or mixed ages. Septic arthritis is more common in children.

What this study adds

- ▶ In this prospective observational study of adults in three EDs in England, leucocyte esterase had a high negative predictive value when evaluating joint fluid for suspected infection.
- ▶ While it cannot distinguish crystal arthropathy from septic joints, this test may help to decrease diagnostic uncertainty and improve adult patient management, safe discharge and flow in the emergency setting. Larger studies are needed.

drug use, rheumatoid arthritis, osteoarthritis, joint surgery, haemodialysis, HIV and immunosuppression.^{4,5}

Proteolytic enzymes which are released by bacteria destroy articular cartilage.⁶ This can occur as early as 1–2 days if left untreated.⁷ A delay to, or suboptimal, treatment can result in significant long-term morbidity or death.^{8,9} Differentiating NJI from other causes of a red, hot, swollen joint can be extremely challenging. Similar presentations are seen with a variety of aetiologies including reactive arthritis, crystalloid arthropathies, haemarthrosis, osteoarthritis, rheumatoid arthritis, lupus, bursitis and trauma. Research by Freed *et al* suggested that it commonly takes up to 3 days to confirm the aetiology on history, examination and synovial examination.¹⁰

Serum white cell count (WCC) and C reactive protein (CRP) along with temperature are useful in the assessment of a patient but these measures lack specificity.^{4,8,11} Currently investigation relies on joint aspiration, visual inspection for pus and analysis with Gram stain while awaiting 48-hour



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Knapper T, Murphy RJ, Rocos B, *et al.* *Emerg Med J* 2021;**38**:707–710.

culture if no organisms are seen on initial microscopy. Gram stain remains the best first-line test at our disposal with 100% specificity for septic arthritis however, it is labour intensive and shows a sensitivity of only 45%.¹² As part of laboratory analysis, a synovial WCC can also be performed. This can be a useful adjunct to Gram staining in differentiating septic from inflammatory conditions, however this is not performed in all laboratories.⁴ Depending on laboratory reference levels, synovial WCC can carry a higher sensitivity than Gram stain but this comes at the cost of specificity.⁴ Unfortunately, with both Gram staining and synovial fluid WCC being laboratory-based investigations, the time period from patient presentation to aspiration to results can be lengthy. In EDs worldwide, with time pressure on management decisions, these laboratory tests often present a barrier to patient flow. In current practice, in the absence of a reliable test, immediate patient management needs to be a clinical decision.

Leucocyte esterase (LE) is an enzyme released by activated leucocytes. Parvizi *et al* analysed its use in investigating prosthetic joint infections, with a positive ($\geq 2+$) LE strip reading yielding a 80.6% sensitivity and 100% specificity.¹³ The utility of LE test strips has also been reported for the analysis of pleural fluid, peritoneal fluid and cerebrospinal fluid to help differentiate between septic and aseptic fluid.^{14–16}

The aim of this multicentre prospective study was to assess the usefulness of LE strip testing in the bedside evaluation of a patient referred with suspected septic arthritis. We hypothesised that LE strip testing of the native joint aspirate could provide a useful adjunct to exclude bacterial septic arthritis and thus allow safe, timely discharge of patients from the ED.

METHODS

A prospective collaborative multicentre observational methodology was used to study the utility of LE in ruling out septic arthritis in three EDs across the Bristol region. The use of this bedside test has been adopted as part of our regional protocol for the assessment of native joint fluid. We performed LE testing on all native joint synovial aspirates in adults (>18 years old) referred to the orthopaedic team from the ED, with suspected septic arthritis at three hospital sites (North Bristol NHS trust, Royal United Hospital NHS foundation trust and Great Western Hospitals NHS foundation trust) between October 2015 and April 2016. At two sites, patient demographics were collected allowing retrospective analysis of blood parameters and admission data. Age, gender, joint, WCC, CRP, duration of symptoms, LE, Gram stain, presence of crystals, 48-hour cultures along with surgical treatment and length of stay were recorded for each case. We excluded any cases of suspected prosthetic joint infection, insufficient aspirate to perform the test and haemarthrosis or blood contamination making the LE result unreadable without further processing.

Joints were aspirated by the on-call orthopaedic doctor under an aseptic technique. Joint aspirates were sent to the microbiology laboratory in a sterile specimen pot for Gram stain, crystals and culture. At the bedside, one drop of the remaining aspirate was then applied to the LE pad on the testing strip, (Combur 7 chemical test strip, Roche diagnostics, Rotkreuz, Switzerland). Results were recorded after 60 s as either neg (white), + (slightly purple), ++ (light purple) or +++ (dark purple) according to the colour chart on the packaging of the test strips as per the manufacturer's instructions. The result of the LE was recorded but was not taken into account in subsequent patient management regarding antibiotics, admission or surgery. An LE test

Table 1 Leucocyte esterase and Gram stain result in relation to the 48-hour culture result

		48-hour culture		
		Positive	Negative	Total
Leucocyte esterase testing	Positive	5	52	57
	Negative	0	23	23
Total		5	75	80
Gram stain	Positive	1	0	1
	Negative	4	75	79
Total		5	75	80

of neg or + (white or slightly purple) was used as a negative indicator. This has previously been shown to have good correlation in the assessment of suspected prosthetic joint infections by Parvizi *et al*.¹³

Sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) with associated 95% CIs were calculated for both LE test and Gram stain using the 48-hour culture results as the gold standard for infection. For the LE test, positive likelihood ratio and negative likelihood ratio were also calculated. No a priori sample size was estimated; the sample size was dependent on the number of patients referred during this time.

Through correlation of admission data with LE and culture results, we estimated the potential cost benefit for a negative LE test result in this study. For this calculation, we used the Department of Health's estimated cost of an acute bed as £303 per day.¹⁷

Patient and public involvement

No patients were involved. LE strip testing formed part of our routine clinical practice prior to conducting this study.

RESULTS

Eighty patients were eligible for inclusion (74% men, 26% women). The cohort had a mean age of 71 years (range 27–96 years). Following data normality testing, mean serum WCC and median CRP were reported for both culture-positive and culture-negative patients. LE test was read as *neg* for nine patients (11%), + for 14 (18%), ++ for 24 (30%) and +++ for 33 patients (41%). Organisms were seen on one Gram stain. Five patients had a positive 48-hour culture (table 1) with pathogens shown in table 2. Infections were seen at all three hospital sites. The five patients were all managed with surgical washout. Of the 52 patients positive for LE but negative on culture, 34 patients were diagnosed with a crystal arthropathy, 17 with a presumed arthritic flare and 1 with no identified cause.

A positive LE result had a sensitivity of 100% (95% CI 47.8% to 100%), specificity 30.7% (95% CI 20.5% to 42.4%), PPV 8.77% (95% CI 7.64% to 10.1%) and positive likelihood ratio

Table 2 Organisms found in positive cultures

Culture-positive organism	No of patients identified with organism
<i>Staphylococcus aureus</i>	1
<i>Pseudomonas aeruginosa</i>	1
Beta haemolytic <i>Streptococcus</i> group C	1
Beta haemolytic <i>Streptococcus</i> group G	1
Beta haemolytic <i>Streptococcus</i> group B	1

Table 3 Sensitivity, specificity, PPV and NPV of leucocyte esterase and Gram stain

Measurement outcome	Result	95% CI
Leucocyte esterase		
Sensitivity (%)	100	47.8 to 100
Specificity (%)	30.7	20.5 to 42.4
PPV (%)	8.77	7.64 to 10.1
NPV (%)	100	82.1 to 100
Gram stain		
Sensitivity (%)	20	0.51 to 71.6
Specificity (%)	100	95.2 to 100
PPV (%)	100	54.6 to 100
NPV (%)	94.9	92.4 to 96.7

NPV, negative predictive value; PPV, positive predictive value.

1.44 (95% CI 1.00 to 1.76) for infection. A negative LE test had an NPV of 100% (95% CI 82.1% to 100%) and negative likelihood ratio 0.00 (95% CI 0.00 to 3.93) in ruling out infection (table 3). There was a 1 in 11 chance of a patient with an LE reading of 2+ or 3+ having an NJI but there were no incidences of a patient with an NJI having a negative or 1+ LE result. In comparison, Gram stain had a sensitivity of 20% (95% CI 0.51% to 71.6%), specificity 100% (95% CI 95.2% to 100%), PPV 100% (95% CI 54.6% to 100%), NPV of 94.9% (95% CI 92.4% to 96.7%) and negative likelihood ratio 0.80 (95% CI 0.52 to 1.24) (table 3).

Clinical and demographic data were collected for 53 patients at two EDs (table 4). Of these, 34 were admitted. A final diagnosis of non-infective pathology was made in 30 of these admissions. Admissions for culture-negative patients accounted for 311 bed-days with a median stay of 6 days (range 1–72 days). The large range related to two admissions where non-orthopaedic infections (urosepsis and pneumonia) were diagnosed and treated during their in-hospital stay. Of the 311 bed-days, 57 days were made up of patients who had a negative LE test result on admission. If admission could have been prevented on the basis of a negative bedside LE test, potential savings of £17271 could have been achieved across these two trusts.

DISCUSSION

Acute presentations of adult patients with a hot, inflamed native joint to the ED are frequent, however, the majority of atraumatic swellings are secondary to non-infective causes.¹ Currently there is no ideal, widely accepted immediate bedside test or marker

Table 4 Comparing the demography, WCC, CRP and incidence of crystal arthropathy for patients with culture-positive and culture-negative results

	Cohort—hospital sites one and two N=53	
	Culture positive	Culture negative
Number of patients	4	49
Number of male	2 (50%)	37 (75%)
Age	81 (75–85)	67.4 (27–96)
Admissions	4/4 (100%)	30/49 (61.2%)
WCC (10 ⁹ /L) mean (SD)	11.4 (2.18)	11.5 (3.80)
CRP (mg/L) median (IQR)	165 (91–434)	71 (43–126)
Crystal arthropathy	0	34

CRP, C reactive protein; WCC, white cell count.

to differentiate infectious from non-infectious joint swellings. Among current diagnostics, peripheral WCCs are raised in only 50% of cases with sensitivity ranging from 23% to 75%.^{4,8} CRP and erythrocyte sedimentation rate (ESR) are acute phase reactants that respond to both infection and inflammation; specificity is low despite their high sensitivity. Literature suggests an ESR >30 mm/hour carries a sensitivity of 76%–97%, however specificity is only 29% for NJI in adults.¹¹ Similarly, CRP values of >100 mg/L have a reported sensitivity around 80% but specificity ranging from 27% to 70%.¹¹ Laboratory Gram stains are universally performed due to their high specificity but sensitivity remains low. A recent study of 830 native joint aspirates for suspected NJI demonstrated sensitivity of 17% (95% CI 10.2% to 25.8%).¹⁸

An accurate rapid diagnosis of NJI is only possible with a positive Gram stain or aspiration of pus. Otherwise, patients presenting with a suspected NJI may be obliged to wait for extended periods in the ED or be admitted for observation or empirical treatment. An audit by Eid *et al* of 60 patients presenting with atraumatic knee effusions reported 24 admissions for empirical antibiotic therapy following aspiration. Only four cases were confirmed septic arthritis. Median stay for all admissions was 4 days (range 2–14 days) with seven medical complications during admission.¹⁹

In our study, we identified a total of 57 orthopaedic bed-days occupied by patients where NJI was excluded by a negative LE test. Discharging these patients could have achieved potential savings of £17271 across two of the trusts in this cohort.¹⁷ However, we appreciate that some of the LE-negative patients may have had concomitant medical or social factors that necessitate admission. The ability to exclude septic arthritis at the point of access could enable earlier assessment, management or admission under the appropriate specialty for the patient's care needs.

LE dipstick testing is a widely adopted, cheap and readily available test as part of the assessment of joint fluid in suspected prosthetic joint infections. To date, the evidence surrounding its use in the assessment of native joint fluid is limited. In this study, LE showed excellent sensitivity with an NPV of 100% (95% CI 82.1% to 100%). These results are in keeping with two recent papers both reporting a 100% NPV for LE.^{20,21} However, we found specificity was poor, 30.7% (95% CI 20.5% to 42.4%) and a PPV of only 8.77% (95% CI 7.64% to 10.1%). Colvin *et al* analysed synovial fluid from five patients with suspected native joint infection reporting similarly low specificity and PPV of 50% and 33%, respectively.²⁰ In contrast, Gautam *et al* reported high specificity, 83%, and PPV, 95%.²¹ The higher specificity and PPV reported by Gautam *et al* may be explained by the predominantly paediatric cohort of patients in their study, over 75% of the patients were under 20 years old. One would expect a different performance of the LE test due to the higher incidence of septic arthritis and rare occurrence of crystal arthropathy in this population group. This increased specificity and PPV of LE for septic arthritis in a paediatric cohort has also been demonstrated by Mortazavi *et al*.²²

The inflammatory response seen with crystal arthropathies means that a positive LE test alone cannot differentiate between NJI and crystal arthropathy. Omar *et al* hypothesised that when dipstick synovial glucose readings were taken alongside the LE test, this would increase the ability of the dipstick to detect septic arthritis.²³ Preliminary results have shown sensitivity 89.5%, specificity 99.2%, PPV 94.4% and NPV 98.4% for a positive LE test (++ or +++) with negative glucose reading for the diagnosis of septic arthritis.²³ Combining the results of LE and glucose dipstick testing may allow distinction of inflammatory

arthropathy from NJI, adding to the usefulness of bedside LE testing.

We acknowledge several limitations of this present study. Although data were collected across three centres, the low incidence of NJI in the adult population and low prevalence in patients referred with a possible NJI meant we had few cases of true NJI. The low prevalence of this disease has the potential to impact on the accuracy of predictive values: increasing the observed NPV and decreasing the PPV. As a multicentre study, there were several doctors performing and interpreting the result of the LE. These doctors were not blinded to the patients' history, examination findings or admission bloods which may have introduced an element of bias. Although this study was performed in a solely adult population, the applicability of an LE to the paediatric population has been demonstrated by Mortazavi *et al*^{21 22}

We would recommend further multicentre investigation to increase numbers and corroborate these results. Using the data from this study as pilot data, assuming a prevalence of NJI of 6.25% (95% CI 2.7% to 13.8%), to test the assumption that the sensitivity of LE testing for detecting NJI is 95%±5%, a sample size of 1138 would be required.²⁴

This study supports the use of bedside LE testing of synovial fluid aspirate in cases of suspected native joint septic arthritis. The test has a high NPV and can act as an adjunct in the decision-making process to help support safe discharge of patients with negative results. These discharged patients may then be followed up with formal laboratory culture results in an outpatient, telephone or primary care setting at 48–72 hours. The LE test can reduce both diagnostic uncertainty and costs to the healthcare system.

Contributors All authors listed on the manuscript contributed substantially to both the project and production of the manuscript. TK, RJM and BR wrote the protocol with supervision from MRW. TK, RJM and BR were joint study leads at one site and NM and JF were the study leads at the other sites. All authors were responsible for data collection and processing. MRW wrote the data analysis and power calculations. All authors were involved in the writing and editing process of the manuscript and all authors have approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval In the analysis of the usefulness of leucocyte esterase, there was no change or impact on patient care or management therefore ethical approval was not required. This was confirmed using the Health Research Authority decision tool.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplemental information.

ORCID iDs

Thomas Knapper <http://orcid.org/0000-0003-0408-1189>

Richard J Murphy <http://orcid.org/0000-0002-5088-8525>

Michael Richard Whitehouse <http://orcid.org/0000-0003-2436-9024>

REFERENCES

- Rutherford AI, Subesinghe S, Bharucha T, *et al*. A population study of the reported incidence of native joint septic arthritis in the United Kingdom between 1998 and 2013. *Rheumatology* 2016;55:2176–80.
- Mathews CJ, Coakley G. Septic arthritis: current diagnostic and therapeutic algorithm. *Curr Opin Rheumatol* 2008;20:457–62.
- Kaandorp CJ, Van Schaardenburg D, Krijnen P, *et al*. Risk factors for septic arthritis in patients with joint disease. A prospective study. *Arthritis Rheum* 1995;38:1819–25.
- Margaretten ME, Kohlswes J, Moore D, *et al*. Does this adult patient have septic arthritis? *JAMA* 2007;297:1478–88.
- Morrey B, Sotelo S, Morrey M. *Morrey's the elbow and its disorders*. 5th ed. Philadelphia, PA: Elsevier, 2017: 756–9.
- Shirliff ME, Mader JT. Acute septic arthritis. *Clin Microbiol Rev* 2002;15:527–44.
- Riegels-Nielson P, Frimodt-Møller N, Jensen JS. Rabbit model of septic arthritis. *Acta Orthop Scand* 1987;58:14–19.
- Weston VC, Jones AC, Bradbury N, *et al*. Clinical features and outcome of septic arthritis in a single UK health district 1982–1991. *Ann Rheum Dis* 1999;58:214–9.
- Söderquist B, Jones I, Fredlund H, *et al*. Bacterial or crystal-associated arthritis? Discriminating ability of serum inflammatory markers. *Scand J Infect Dis* 1998;30:591–6.
- Freed JF, Nies KM, Boyer RS. Acute monoarticular arthritis. *JAMA* 1980;243:2314–6.
- Carpenter CR, Schuur JD, Everett WW, *et al*. Evidence-based diagnostics: adult septic arthritis. *Acad Emerg Med* 2011;18:781–96.
- Faraj AA, Omonbude OD, Godwin P. Gram staining in the diagnosis of acute septic arthritis. *Acta Orthop Belg* 2002;68:388–91.
- Parvizi J, Jacovides C, Antoci V, *et al*. Diagnosis of periprosthetic joint infection: the utility of a simple yet unappreciated enzyme. *J Bone Joint Surg Am* 2011;93:2242–8.
- Azoulay E, Fartoukh M, Galliot R, *et al*. Rapid diagnosis of infectious pleural effusions by use of reagent strips. *Clin Infect Dis* 2000;31:914–9.
- Bortosh W, Siedner M, Carroll RW. Utility of the urine reagent strip leucocyte esterase assay for the diagnosis of meningitis in resource-limited settings: meta-analysis. *Trop Med Int Health* 2017;22:1072–80.
- Castellote J, López C, Gornals J, *et al*. Rapid diagnosis of spontaneous bacterial peritonitis by use of reagent strips. *Hepatology* 2003;37:893–6.
- Department of Health. Reference costs 2014–15, 2015. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/477919/2014-15_Reference_costs_publication.pdf
- Gbejuade H, Elsakka M, Cutler L. How well does synovial fluid gram staining correlate with cultures in native joint infections? *Orthop Rev* 2019;11:8156.
- Eid A, Burrows V, Murray J. Are we managing knee effusions well? *British Journal of Medical Practitioners* 2012;5:a504.
- Colvin OC, Kransdorf MJ, Roberts CC, *et al*. Leukocyte esterase analysis in the diagnosis of joint infection: can we make a diagnosis using a simple urine dipstick? *Skeletal Radiol* 2015;44:673–7.
- Gautam VK, Saini R, Sharma S. Effectiveness of leucocyte esterase as a diagnostic test for acute septic arthritis. *J Orthop Surg* 2017;25:230949901668501–6.
- Mortazavi SMJ, Kalantar H, Baghdadi S, *et al*. The utility of leucocyte esterase strip test in the diagnosis of pediatric septic arthritis. *J Pediatr Orthop* 2020;40:e312–6.
- Omar M, Ettinger M, Reichling M, *et al*. Preliminary results of a new test for rapid diagnosis of septic arthritis with use of leucocyte esterase and glucose reagent strips. *J Bone Joint Surg Am* 2014;96:2032–7.
- Carley S, Dosman S, Jones SR, *et al*. Simple nomograms to calculate sample size in diagnostic studies. *Emerg Med J* 2005;22:180–1.